arXiv:hep-lat/0312041v1 29 Dec 2003

A RELATIVISTIC INVARIANT SCHEME FOR THE QUANTUM KLEIN-GORDON AND DIRAC FIELDS ON THE LATTICE

Miguel Lorente. *Departamento de Física, Universidad de Oviedo, 33007 Oviedo, Spain*

Abstract. We explore the connection between the symmetry transformations and conservation laws for the Klein-Gordon and Dirac fields on the lattice. The generators of the space time translations and Lorentz boost (defined on the lattice) are constants of motion and satisfy the standard commutation relations.

1. Quantization of the Klein-Gordon field

We introduce the method of finite differences for the Klein-Gordon scalar field. A scheme for the wave equation consistent with the continuous case (the truncation error is of second orden with respect to space and time variables) can be constructed as follows:

$$\left(\frac{1}{\tau^2}\nabla_n\triangle_n\tilde{\nabla}_j\tilde{\triangle}_j-\frac{1}{\varepsilon^2}\nabla_j\triangle_j\tilde{\nabla}_n\tilde{\triangle}_n+M^2\tilde{\nabla}_n\tilde{\triangle}_n\tilde{\nabla}_j\tilde{\triangle}_j\right)\phi_j^n=0 \tag{1.1}$$

where the field is defined in the grid points of the (1+1)-dimensional lattice $\phi_j^n \equiv \phi\,(j\varepsilon, n\tau)$, ε, τ being the space and time fundamental intervals, j, n integer numbers and

$$\Delta f_j \equiv f_{j+1}-f_j \quad , \quad \nabla f_j = f_j - f_{j-1}$$
$$\tilde{\Delta} f_j \equiv \frac{1}{2}(f_{j+1}+f_j) \quad , \quad \tilde{\nabla} f_j \equiv \frac{1}{2}(f_j+f_{j-1})$$

and similarly for the time index.

Using the method of separation of variables it can easily be proved that the following functions of discrete variables are solutions of the wave equation (1.1):

$$f_j^n\,(k,\omega)=\left(\frac{1+\frac{1}{2}i\varepsilon k}{1-\frac{1}{2}i\varepsilon k}\right)^j\left(\frac{1-\frac{1}{2}i\tau\omega}{1+\frac{1}{2}i\tau\omega}\right)^n \tag{1.2}$$

provided the “dispersion relation is satisfied: $\omega^2 - k^2 = M^2$ (1.3)
M being the mass of the particle.

In the limit, $j \to \infty$, $n \to \infty$, $j\varepsilon \to x$, $n\tau \to t$ the functions (1.2) becomes plane wave solutions

$$f_j^n(k,\omega) \to exp\ i(kx-\omega t) \tag{1.4}$$

Imposing boundary conditions on the space indices

$$f_0^n(k,\omega) = f_N^n(k,\omega) \tag{1.5}$$

we get $k_m = \frac{2}{\varepsilon} tg \frac{\pi m}{N}$, $m = 0, 1, \ldots, N-1$ (1.6)

For the positive energy solutions we define $\omega_m = +(k_m^2 + M^2)^{\frac{1}{2}}$ (1.7)

For $n = 0$ the functions $f_j(k_m)$ satisfy the orthogonality relations

$$\frac{1}{N} \sum_{j=0}^{N-1} f_j^*(k_m) f_j(k_{m'}) = \delta_{mm'} \tag{1.8}$$

For the quantization of the Klein-Gordon field we impose the equal time commutations relations:

$$[\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n, \tilde{\nabla}_{j'} \tilde{\Delta}_{j'} \pi_{j'}^n] = \frac{i}{\varepsilon} \delta_{jj'} \tag{1.12}$$

$$[\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n, \tilde{\nabla}_{j'} \tilde{\Delta}_{j'} \phi_{j'}^n] = 0 = [\tilde{\nabla}_j \tilde{\Delta}_j \pi_j^n, \tilde{\nabla}_{k'} \tilde{\Delta}_{j'} \pi_{j'}^n] \tag{1.13}$$

The Hamiltonian for the real scalar field reads:

$$\mathrm{H}_n = \varepsilon \sum_{j=0}^{N-1} \frac{1}{2} \left\{ \left(\tilde{\nabla}_j \tilde{\Delta}_j \pi_j^n \right)^2 - \frac{1}{\varepsilon^2} \left(\nabla_j \Delta_j \phi_j^n \right) \left(\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n \right) + M^2 \left(\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n \right)^2 \right\} \tag{1.14}$$

A convenient scheme for the Heisenberg equations of motion is

$$\frac{1}{\tau} \Delta_n \left(\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n \right) = \frac{1}{i} \left[\tilde{\Delta}_n \left(\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n \right), H_n \right] \tag{1.15}$$

$$\frac{1}{\tau} \Delta_n \left(\tilde{\nabla}_j \tilde{\Delta}_j \pi_j^n \right) = \frac{1}{i} \left[\tilde{\Delta}_n \left(\tilde{\nabla}_j \tilde{\Delta}_j \pi_j^n \right), H_n \right] \tag{1.16}$$

Using (1.12) and (1.13) we get

$$\frac{1}{\tau} \Delta_n \left(\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n \right) = \frac{1}{i} \tilde{\Delta}_n \left[\tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n, H_n \right] = \Delta_n \left(\tilde{\nabla}_j \tilde{\Delta}_j \pi_j^n \right) \tag{1.17}$$

$$\frac{1}{\tau} \Delta_n \left(\tilde{\nabla}_j \tilde{\Delta}_j \pi_j^n \right) = \Delta_n \left\{ \frac{1}{\varepsilon^2} \nabla_j \Delta_j \phi_j^n - M^2 \tilde{\nabla}_j \tilde{\Delta}_j \phi_j^n \right\} \tag{1.18}$$

leading to the wave equation (1.1) for the fields ϕ_j^n and π_j^n.

Since the plane wave solutions $f_j^n(k_m, \omega_m)$, $(m = 0, 1, \ldots, N-1)$ form a complete set of orthogonal functions with respect to the space indices, we can expand the wave field and its conjugate momentum as

$$\phi_j^n = \frac{1}{\sqrt{N\varepsilon}} \sum_{m=-N/2}^{N/2-1} \frac{\left(1 + \frac{1}{4}\varepsilon^2 k_m\right)}{\sqrt{2\omega_m}} \Big(a_m f_j^n \left(k_m, \omega_m\right) + a_m^\dagger f_j^{*n} \left(k_m, \omega_m\right) \Big) \tag{1.19}$$

$$\pi_j^n = \frac{-i}{\sqrt{N\varepsilon}} \sum_{m=-N/2}^{N/2-1} \sqrt{\frac{\omega_m}{2}} \left(1 + \frac{1}{4}\varepsilon^2 k_m \right) \Big(a_m f_j^n \left(k_m, \omega_m\right) - a_m^\dagger f_j^{*n} \left(k_m, \omega_m\right) \Big) \tag{1.20}$$

Inverting this equations we can calculate with the help of (1.12) and (1.13):

$$[a_m, a_{m'}^\dagger] = \delta_{mm'} \quad , \quad [a_m, a_{m'}] = 0 \quad , \quad [a_m^\dagger, a_{m'}^\dagger] = 0 \tag{1.21}$$

Substituting (1.19) and (1.20) in (1.14) we find

$$\mathrm{H} = \frac{1}{2} \sum_{\mathrm{m}=-\mathrm{N}/2}^{\mathrm{N}/2-1} \omega_{\mathrm{m}} \left(\mathrm{a_m a_m^\dagger} + \mathrm{a_m^\dagger a_m} \right) \tag{1.22}$$

If we define the linear momentum on the lattice as

$$P = - \sum_{j=0}^{N-1} \left(\tilde{\nabla}_j \tilde{\Delta}_j \pi_j^n \right) \left(\tilde{\nabla}_j \Delta_j \phi_j^n \right) \tag{1.23}$$

we find $P = \sum_{m=-N/2}^{N/2-1} k_m \left(a_m a_m^\dagger + a_m^\dagger a_m \right) = \sum_{m=-N/2}^{N/2-1} k_m a_m^\dagger a_m$ because the zero point momentum vanish by cancellation of k_m with k_{-m}

2. Quantization of the Dirac field

The Hamiltonian for the Dirac field $\psi_{\alpha j}^n$ and its hermitian adjoint $\psi_{\alpha j}^{\dagger n}$ can be defined as:

$$\mathrm{H}_n = \varepsilon \sum_{j=0}^{N-1} \tilde{\Delta}_j \psi_j^{\dagger n} \left\{ \gamma_4 \gamma_1 \frac{1}{\varepsilon} \Delta_j \psi_j^n + M \gamma_4 \tilde{\Delta}_j \psi_j^n \right\} \tag{2.1}$$

with $\gamma_1 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \gamma_4 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, i\gamma_1\gamma_4 = \gamma_5 = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$ (2.2)

We impose the equal time anticommutation relations for the fields $\psi_{\alpha j}^n$ and $\psi_{\alpha j}^{\dagger n}$

$$\left[\tilde{\Delta}_j \psi_{\alpha j}^n, \tilde{\Delta}_{j'} \psi_{\beta j'}^{\dagger n} \right]_+ = \frac{1}{\varepsilon} \delta_{\alpha\beta} \delta_{jj'} \tag{2.3}$$

$$\left[\tilde{\Delta}_j \psi_{\alpha j}^n, \tilde{\Delta}_{j'} \psi_{\beta j'}^n \right]_+ = \left[\tilde{\Delta}_j \psi_{\alpha j}^{\dagger n}, \tilde{\Delta}_{j'} \psi_{\beta j'}^{\dagger n} \right]_+ = 0 \tag{2.4}$$

Using these relations we obtain for the Heisenberg equation of motion

$$\begin{aligned} \frac{1}{\tau} \Delta_n \left(\tilde{\Delta}_j \psi_j^n \right) &= \frac{1}{i} \left[\tilde{\Delta}_n \left(\tilde{\Delta}_j \psi_j^n \right), H_n \right] = \frac{1}{i} \tilde{\Delta}_n \left[\tilde{\Delta}_j \psi_j^n, H_n \right] \\ &= \frac{1}{i} \tilde{\Delta}_n \left\{ \gamma_4 \gamma_1 \frac{1}{\varepsilon} \Delta_j \psi_j^n + M \gamma_4 \tilde{\Delta}_j \psi_j^n \right\} \end{aligned} \tag{2.5}$$

from which it can be derived the Dirac equation on the lattice:

$$\left(\gamma_1 \frac{1}{\varepsilon} \Delta_j \tilde{\Delta}_n - i \gamma_4 \frac{1}{\tau} \Delta_n \tilde{\Delta}_j + M \tilde{\Delta}_j \tilde{\Delta}_n \right) \psi_j^n = 0 \tag{2.6}$$

Applying the operator $\gamma_1 \frac{1}{\varepsilon} \nabla_j \tilde{\nabla}_n - i \gamma_4 \frac{1}{\tau} \nabla_n \tilde{\nabla}_j - M \tilde{\nabla}_j \tilde{\nabla}_n$ on both sides of (2.6) we recover the wave equation (1.1) for ψ_j^n.

3. Conservation laws and Lorentz invariance

As in the continuous case we can make the connection between symmetries and conservation laws in the language of generators. The condition for symmetry under u space and time displacement and pure Lorentz transformation is that the generators are constant of the motion . In the case of the Klein-Gordon fields the generators of the (one step) space and time translations and Lorentz boost can be taken as:

$$P=-\sum_{j=0}^{N-1}\frac{1}{2}\left\{\left(\tilde{\nabla}_j\tilde{\Delta}_j\pi_j^n\right)\left(\tilde{\nabla}_j\Delta_j\phi_j^n\right)+\left(\tilde{\nabla}_j\Delta_j\phi_j^n\right)\left(\tilde{\nabla}_j\tilde{\Delta}_j\pi_j^n\right)\right\} \tag{3.1}$$

$$H=\varepsilon\sum_{j=0}^{N-1}\frac{1}{2}\left\{\left(\tilde{\nabla}_j\tilde{\Delta}_j\pi_j^n\right)^2+\frac{1}{\varepsilon^2}\left(\tilde{\nabla}_j\Delta_j\phi_j^n\right)^2+M^2\left(\tilde{\nabla}_j\tilde{\Delta}_j\phi_j^n\right)^2\right\} \tag{3.2}$$

$$\begin{aligned}K=&\,\varepsilon\sum_{j=0}^{N-1}\frac{1}{2}\varepsilon j\left\{\left(\tilde{\Delta}_j\tilde{\nabla}_j\tilde{\Delta}_j\pi_j^n\right)^2+\frac{1}{\varepsilon^2}\left(\tilde{\Delta}_j\tilde{\nabla}_j\Delta_j\phi_j^n\right)^2+M^2\left(\tilde{\Delta}_j\tilde{\nabla}_j\tilde{\Delta}_j\phi_j^n\right)^2\right\}\\&-\varepsilon\sum_{j=0}^{N-1}n\tau\frac{1}{2}\left\{\left(\tilde{\Delta}_j\tilde{\nabla}_j\phi_j^n\right)\left(\tilde{\Delta}_j\nabla_j\phi_j^n\right)+\left(\tilde{\Delta}_j\nabla_j\phi_j^n\right)\left(\tilde{\Delta}_j\tilde{\nabla}_j\pi_j^n\right)\right\}\end{aligned} \tag{3.3}$$

Using (1.17) and (1.18) it can be proved that these operators are constant of time

$$\frac{1}{\tau}\Delta_nP=\frac{1}{\tau}\Delta_nH=\frac{1}{\tau}\Delta_nK=0 \tag{3.4}$$

In order to check the Lorentz invariance of quantized field scheme on the lattice, one can prove with the help of (1.12) and (1.13) that these operators satisfy the standard commutation relations:

$$[H,P]=0\qquad,\qquad[K,H]=iP\qquad,\qquad[K,P]=iH \tag{3.5}$$

For the Dirac quantum fields, the generators of the (one step) space and time translations and Lorentz boost can be taken as

$$P=-i\sum_{j=0}^{N-1}(\tilde{\Delta}_j\psi_{\alpha j}^{\dagger n})(\Delta_j\psi_{\alpha j}^n) \tag{3.6}$$

$$H=\varepsilon\sum_{j=0}^{N-1}\tilde{\Delta}_j\psi_{\alpha j}^{\dagger n}\{(\gamma_4\gamma_1)_{\alpha\beta}\frac{1}{\varepsilon}\Delta_j\psi_{\beta j}^n+M(\gamma_4)_{\alpha\beta}\tilde{\Delta}_j\psi_{\beta j}^n\} \tag{3.7}$$

$$\begin{aligned}M_{14}=&\,i\varepsilon\sum_{j=0}^{N-1}\varepsilon j\{\tilde{\Delta}_j\tilde{\nabla}_j\psi_{\alpha j}^{\dagger n}\{(\gamma_4\gamma_1)_{\alpha\beta}\frac{1}{\varepsilon}\Delta_j\tilde{\nabla}_j\psi_{\beta j}^n+M(\gamma_4)_{\alpha\beta}\tilde{\Delta}_j\tilde{\nabla}_j\psi_{\beta j}^n\}\}\\&-i\varepsilon\sum_{j=0}^{N-1}\tau n\{\tilde{\Delta}_j\psi_{\alpha j}^{\dagger n}\Delta_j\psi_{\alpha j}^n\}+\varepsilon\sum_{j=0}^{N-1}\tilde{\Delta}\psi_{\alpha j}^{\dagger}\frac{1}{2i}(\gamma_1\gamma_4)_{\alpha\beta}\tilde{\Delta}_j\psi_{\beta j}^n\end{aligned} \tag{3.8}$$

Using (2.5), (2,3) and (2,4) it can be proved that these operators are constant of time

$$\frac{1}{\tau}\Delta_nP=\frac{1}{\tau}\Delta_nH=\frac{1}{\tau}\Delta_nM_{14}=0 \tag{3.9}$$

and that they satisfy the standard commutation relations

$$[H,P]=0\quad,\quad[H,M_{14}]=P\quad,\quad[P,M_{14}]=H \tag{3.10}$$